

Lower bounds for average sample size and efficiency of sequential selection procedures

Kareev I.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

In this paper we establish lower bounds for average sample size in procedures for selecting a population with the largest value of scalar parameter, when the given probability of the correct solution is guaranteed. In construction of bounds we use universal bounds of Volodin and Malyutov [I. N. Volodin, Theory Probab. Appl., 24 (1979), pp. 120-129], [M. B. Malyutov, Izv. Vyssh. Uchebn. Zaved. Mat., 1983, no. 11, pp. 19-41 (in Russian)]. The obtained results are applied to the estimation of efficiency of the Bechhofer selection procedure [R. E. Bechhofer, Ann. Math. Statist., 25 (1954), pp. 16-39] and sequential modification of it [S. C. Kao and T. L. Lai, Comm. Statist. A-Theory Methods, 9 (1980), pp. 1657-1676], [R. E. Bechhofer, J. Kiefer, and M. Sobel, Sequential Identification and Ranking Procedures, with Special Reference to Koopman-Darmois Populations, University of Chicago Press, Chicago, 1968]. ©2013 Society for Industrial and Applied Mathematics.

<http://dx.doi.org/10.1137/S0040585X97985935>

Keywords

Efficiency, Lower bounds for average sample size, Selection